

COLOR-FLAVOR (UN-) LOCKING

M. OERTEL¹ and M. BUBALLA^{2,3}

¹ *IPN Lyon; 43, bv du 11 novembre 1918; F-69622 Villeurbanne Cédex, France*

² *IKP, TU Darmstadt, Schlossgartenstr. 9, D-64289 Darmstadt, Germany*

³ *GSI Darmstadt; Postfach 110552; D-64220 Darmstadt; Germany*

Abstract

The structure of the phase diagram of strongly interacting matter at moderate densities is calculated within a 3-flavor NJL-type quark model with realistic quark masses. We focus on the influence of the selfconsistently determined effective strange quark mass on the color-flavor unlocking phase transition.

1 Introduction

The structure of the QCD phase diagram as a function of temperature and density has been studied extensively throughout the last years, on theoretical as well as on experimental grounds. For a long time the discussion has been restricted to two phases: the hadronic phase and the quark-gluon-plasma. This has dramatically changed after the discovery that, already for densities present in the interior of neutron stars, color superconducting phases might exist with energy gaps of the order of $\Delta \sim 100$ MeV [1].

At these moderate densities QCD cannot be treated perturbatively and therefore one has to rely on model calculations. In general local two-quark interactions are chosen which are motivated, e.g., from instanton interactions. Most interactions favor a condensation in the scalar color- $\bar{3}$ channel:

$$s_{AA'} = \langle \psi^T C \gamma_5 \tau_A \lambda_{A'} \psi \rangle . \quad (1)$$

Here both, τ_A and $\lambda_{A'}$ are the antisymmetric generators of $SU(3)$, i.e., the antisymmetric Gell-Mann matrices ($A, A' \in \{2, 5, 7\}$), acting in flavor and color space, respectively.

Let us first discuss two limiting cases: Assuming the mass of the strange quark to be infinite, we can neglect pairing between strange and lighter quarks and the flavor index in Eq. (1) is restricted to $A = 2$. The three condensates $s_{2A'}$ form a vector in color space, which always can be rotated into the $A' = 2$ -direction. Hence the two-flavor superconducting state (2SC) state can be

characterized by $s_{22} \neq 0$ and $s_{AA'} = 0$ if $(A, A') \neq (2, 2)$. In this state color $SU(3)$ is spontaneously broken down to $SU(2)$. For massless u and d quarks the 2SC state is invariant under chiral $SU(2)_L \times SU(2)_R$ transformations.

The second limiting case is to assume three degenerate quark flavors. Then dense matter is expected to form a so-called color-flavor locked (CFL) state [2], characterized by the situation $s_{22} = s_{55} = s_{77} \neq 0$ and $s_{AA'} = 0$ if $A \neq A'$. In this state $SU(3)_{color} \times SU(3)_L \times SU(3)_R$ and the $U(1)$ -symmetry related to baryon-number conservation are broken down to a common $SU(3)_{color+V}$ subgroup where color and flavor rotations are locked.

Both situations discussed above are idealizations of the real world. For sufficiently large quark chemical potentials $\mu \gg M_s$, the s quark mass becomes of course almost negligible against μ and matter is expected to be in the CFL phase. It is not clear, however, whether this CFL phase is directly connected to the hadronic phase at low-densities, or whether an intermediate 2SC phase exist, where only up and down quarks are paired. It is obvious that the answer to this question depends on the strange quark mass which is in general T - and μ -dependent. Moreover, it could depend on the presence of quark-antiquark and diquark condensates. This means, not only the phase structure depends on the quark mass, but also the quark mass depends on the phase.

To study these interdependencies we will use an NJL-type model, where the masses are closely related to the (T - and μ -dependent) $\bar{q}q$ condensates [3]

$$\phi_u = \langle \bar{u}u \rangle = \langle \bar{d}d \rangle \quad \text{and} \quad \phi_s = \langle \bar{s}s \rangle. \quad (2)$$

This work is largely based on Ref. [4]. In addition we will discuss the effects of a flavor mixing interaction, which has not been included in that reference.

2 Model

We consider the effective Lagrangian

$$\mathcal{L}_{eff} = \bar{\psi}(i\partial - \hat{m})\psi + \mathcal{L}_{q\bar{q}} + \mathcal{L}_{qq}, \quad (3)$$

where ψ denotes a quark field with three flavors and three colors. The mass matrix $\hat{m}$ has the form $\hat{m} = \text{diag}(m_u, m_d, m_s)$ in flavor space. Throughout this article we assume isospin symmetry, $m_u = m_d$.

To study the interplay between the color-superconducting diquark condensates $s_{AA'}$ and the quark-antiquark condensates ϕ_u and ϕ_s we consider an NJL-type interaction with a quark-quark part

$$\mathcal{L}_{qq} = H \sum_{A=2,5,7} \sum_{A'=2,5,7} (\bar{\psi} i\gamma_5 \tau_A \lambda_{A'} C \bar{\psi}^T)(\psi^T C i\gamma_5 \tau_A \lambda_{A'} \psi). \quad (4)$$

and a quark-antiquark part

$$\begin{aligned} \mathcal{L}_{q\bar{q}} = & G \left\{ \sum_{a=1}^8 \left[(\bar{\psi} \tau_a \psi)^2 + (\bar{\psi} i \gamma_5 \tau_a \psi)^2 \right] + \frac{2}{3} \left[(\bar{\psi} \psi)^2 + (\bar{\psi} i \gamma_5 \psi)^2 \right] \right\} \\ & - K \left[\det_f \left(\bar{\psi} (1 + \gamma_5) \psi \right) + \det_f \left(\bar{\psi} (1 - \gamma_5) \psi \right) \right]. \end{aligned} \quad (5)$$

The latter consists of a $U(3)_L \times U(3)_R$ -symmetric 4-point interaction and a 't Hooft-type 6-point interaction which breaks the $U_A(1)$ symmetry.

Starting from this Lagrangian we calculate the mean-field thermodynamic potential Ω at temperature T and quark chemical potential μ and determine the selfconsistent solutions for the expectation values $\phi_u = \phi_d$, ϕ_s , s_{22} , and $s_{55} = s_{77}$ by minimizing Ω with respect to these expectation values. In this context it is convenient to introduce the effective quark masses $M_i = m_i - 4G\phi_i + 2K\phi_j\phi_k$, $((i, j, k) = \text{cyclic permutations of } (u, d, s))$ and the diquark gaps $\Delta_i = -2Hs_{ii}$.

3 Numerical results

To determine the values of the various condensates we first have to specify the parameters of the interaction. As a starting point we take the parameter values obtained in Ref. [5] by fitting vacuum masses and decay constants of pseudoscalar mesons. It is, however, certainly worthwhile to study in detail the dependence of the results on the strength of the 't Hooft-type interaction which mixes different flavors and thus affects M_s even in regimes, where no strange quark states are populated. To that end we vary the coupling constant K and readjust G and m_s such that the vacuum masses stay unchanged. Since the quark-quark part of the interaction does not influence any vacuum properties the coupling constant H has to be determined in a different way. For $K = 0$ we take $H = 3/4G$ which corresponds to the ratio of G and H obtained from a four-fermion interaction with the quantum numbers of a single-gluon exchange [2, 6]. The resulting value for H is then kept constant for all values of K . As in Ref. [5], all quark loops are regularized by a sharp 3-momentum cut-off Λ .

We begin with the discussion of the results at $T = 0$. The behavior of the constituent quark masses and the diquark gaps as functions of μ is displayed in Fig. 1. The left panel corresponds to $K = 0$, the right panel to $K\Lambda^5 = 20$. One can clearly distinguish three phases. At low μ , the diquark gaps vanish and the constituent quark masses stay at their vacuum values. Hence, in a very schematic sense, this phase can be identified with the ‘‘hadronic phase’’.




Figure 1: Gap parameters at $T = 0$ as functions of μ : Constituent masses $M_u = M_d$ (solid) and M_s (dashed), and diquark gaps Δ_2 (dotted) and Δ_5 (dashed-dotted). Left: $K = 0$. Right: $K\Lambda^5 = 20$.

At a critical chemical potential $\mu = \mu_1$ a first-order phase transition to the 2SC phase takes place: The diquark gap Δ_2 has now a non-vanishing value, whereas Δ_5 remains zero. At the same time the mass of the light quarks drops from the vacuum value to about 50 MeV and the baryon number density jumps from zero to about 2.5 nuclear matter density. This behavior, including the value of μ_1 , is rather independent of K . Obviously this is not the case for M_s : For $K = 0$ the strange quarks are completely unaffected, whereas a nonzero value of K leads to a considerably lower strange quark mass in the 2SC phase. This favors the population of strange quark states already in the 2SC phase and indeed we observe a nonzero density of strange quarks above $\mu \approx 415$ MeV. This is the reason for the accelerated decrease of M_s which can be observed in Fig. 1.

At $\mu = \mu_2$ the system undergoes a second first-order phase transition, this time from the 2SC phase into the CFL phase, which is characterized by a non-vanishing diquark gap Δ_5 (together with a non-vanishing Δ_2). The flavor-mixing interaction accelerates the transition to the CFL phase: The value of μ_2 for $K\Lambda^5 = 20$ is about 50 MeV lower than for $K = 0$. On the other hand, the values of Δ_5 and Δ_2 for a given value of μ depend only weakly on K , as long as the CFL solution is stable. Since in both cases there still is a metastable CFL solution below μ_2 , the main effect of the flavor mixing interaction is to render the CFL solution stable for lower values of μ .

We now extend our analysis to non-vanishing temperatures. The resulting phase diagrams in the μ - T plane are shown in Fig. 2. The left panel corresponds to the results of Ref. [4], obtained with a slightly different parameter set with $K = 0$. We can distinguish four different regimes: At low temperatures we find




Figure 2: Left: Phase diagram for $K = 0$ (from [4]). The lines indicate phase boundaries with first-order (solid) and second-order (dashed) phase transitions. Right: Phase diagram for $K\Lambda^5 = 5$ (dashed), 12.36 (dotted), and 20 (solid).

(with increasing μ) the “hadronic phase”, the 2SC phase, and the CFL phase. Similar to zero temperature these three phases are well separated by first-order phase transitions. The high temperature regime is governed by the QGP phase, which is characterized by vanishing diquark condensates and small values of ϕ_u and (for large enough μ) ϕ_s . There we find smooth crossovers with respect to ϕ_u and ϕ_s instead of the first-order phase transitions, which therefore end in second order endpoints. The transition from the 2SC phase to the QGP phase is of second order and the critical temperature is in almost perfect agreement with the well-known BCS relation $T_c = 0.57\Delta_2(T=0)$.

Since the diquark gap Δ_5 vanishes at a somewhat lower temperature than Δ_2 , there is an intermediate 2SC phase “above” the CFL phase. It has been argued [6] that this color-flavor-unlocking transition has to be first order because pairing between light and strange quarks can only occur if the gap is of the same order as the mismatch between the Fermi surfaces. Hence the value of the gap must be discontinuous at the phase transition. Moreover, the phase transition corresponds to a finite temperature chiral restoration phase transition in a three-flavor theory, and therefore the universality arguments of Ref. [7] should apply [1]. At low temperatures our results are in agreement with these predictions. However, above a critical point we find a second order unlocking transition. In fact, the above arguments are not as stringent as they seem to be on a first sight: First of all the Fermi surfaces are smeared out due to thermal effects and secondly the universality argument is not rigorously applicable because the 2SC phase is not a three-flavor chirally restored phase, but only $SU(2) \times SU(2)$ symmetric. Note that even $SU(3)_V$ is broken explicitly and spontaneously.

On the r.h.s. of Fig. 2 we show first results on the variation of the phase diagram with K . We find that the 't Hooft interaction mainly influences the unlocking transition at low T . The other boundaries remain almost unchanged.

4 Conclusions

We studied the “cold dense” region of strongly interacting matter within the framework of an NJL-type quark model. Our main intention was to examine the effect of selfconsistently determined realistic quark masses, in particular of the strange quark mass. Our results indicate that the 2SC state is the most favored one up to rather high densities. The color-flavor unlocking transition is mostly triggered by a discontinuous change of the strange quark mass. Including a flavor mixing interaction, which lowers the strange quark mass within the 2SC phase, the CFL phase becomes favored for smaller μ . With increasing temperature the values of the diquark condensates decrease and vanish at some critical value T_c which is lower for Δ_5 than for Δ_2 . This color-flavor unlocking transition becomes second order above a critical point.

For simplicity, we restricted our studies to a common baryon chemical potential for all flavors. However, for many applications, like quark cores of neutron stars, it would be interesting to consider β -equilibrated quark matter, where the chemical potential for u quarks is in general different from that of d and s quarks. Moreover, since the effective interaction at the densities of interest is rather unknown, a more systematic examination of the sensitivity of the results on the choice of the interaction should be performed to sort out some common features.

References

- [1] K. Rajagopal and F. Wilczek, hep-ph/0011333, and references therein.
- [2] M. Alford, K. Rajagopal, and F. Wilczek, Nucl. Phys. B 537 (1999) 443.
- [3] T. Hatsuda and T. Kunihiro, Phys. Rep. 247, (1994) 221.
- [4] M. Buballa and M. Oertel, Nucl. Phys. A, in press, hep-ph/0109095.
- [5] P. Rehberg, S.P. Klevansky and J. Hübner, Phys. Rev. C 53 (1996) 410.
- [6] M. Alford, J. Berges, K. Rajagopal, Nucl. Phys. B 558 (1999) 219.
- [7] R.D. Pisarski and F. Wilczek, Phys. Rev. D 29 (1984) 338.